


MAIN FEATURES

- LATEX ENHANCED IMMUNOTURBIDIMETRIC METHOD
- EASY TO PROGRAM ON ANY TYPE OF AUTOMATED ANALYZERS
- GOOD CORRELATION WITH A NEPHELOMETRIC COMMERCIAL PROCEDURE
- RANGE OF ANALYSIS0.0 – 10.0 MG/L
- GOOD PRECISION.....INTER-RUN: CVs < 1.22 %
.....INTRA-RUN: CVs < 3.37 %
- LIQUID REAGENTSREADY TO USE
- SELF-LIFE.....2 YEARS
- PROZONE> 16 MG/L
- LDD< 0.05 MG/L

GENERAL APPLICATION



PRECISION

CONTROL	LEVEL 1	LEVEL 2	LEVEL 3	LEVEL 4
MEAN MG/L	0.86	1.56	3.01	5.03
SD	0.01	0.02	0.03	0.04
CV %	1.02	1.22	1.01	0.70

WITHIN-RUN PRECISION (N=10)

CONTROL	LEVEL 1	LEVEL 2	LEVEL 3	LEVEL 4
MEAN MG/L	0.85	1.53	2.98	5.00
SD	0.03	0.03	0.05	0.14
CV %	3.37	2.18	1.54	2.85

BETWEEN-RUN PRECISION (N=10)

SENSITIVITY*

(* LEAST DETECTABLE DOSE (SENSITIVITY) DETERMINED AS THE CYSTATIN C CONCENTRATION NEEDED FOR A THREE DEVIATION RESPONSE OVER THE 0.0 MG/L STANDARD ONE

CONTROL	LEVEL 1
N	20
LDD MG/L	0.03


SENSITIVITY

ORDER INFORMATION


ANOTHER PRESENTATIONS ARE AVAILABLE ON REQUEST

- REF: K-9590T CYSTATIN C BUFFER 30 ML
- REF: S-9390T CYSTATIN C CALIBRATOR SET 6 x 1 ML
- REF: C-9390T CYSTATIN C CONTROL SET 2 x 1 ML



CALIBRATION CURVE



PROZONE



METHOD COMPARISON



LINEARITY

